

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Medical Paediatrics and Oncology

Journal homepage: <https://www.ijmpo.com/>

Case Report

Cutaneous metastasis in a case of carcinoma floor of mouth- A rare case report

Sheenu Priya¹, Romi Kant Grover^{1*}, Pardeep Garg¹, Navik Goyal¹

¹Dept. of Radiation Oncology, Guru Gobind Singh Medical College and Hospital, Faridkot, Punjab, India



ARTICLE INFO

Article history:

Received 15-12-2023

Accepted 06-02-2024

Available online 26-02-2024

Keywords:

Cutaneous metastasis

Head and neck cancers

Carcinoma larynx

Oral cavity

Skin nodules

ABSTRACT

Background: The incidence of cutaneous metastasis in head and neck cancers is less than 1%. Carcinoma of the oral cavity is one of the most common cancers of the head neck region with frequent metastasis to internal organs, especially the lungs, liver, and bone. We present a case of Carcinoma of floor of mouth with cutaneous metastasis.

Case Presentation: A 55 year old lady presented to our department with complaints of non-healing ulcer in the oral cavity and ulcerated nodules over neck and chest. On examination, mouth opening tongue mobility was restricted with grade 2 trismus. An ulcer in left gb sulcus, bilateral lymphadenopathy and multiple ulcerated skin nodules on neck and chest wall were noted. CECT Face and Neck showed heterogeneously enhancing mass at the floor of mouth involving left buccal mucosa and erosion of mandible, lymph node mets and skin nodules. ECOGPS was 1. She was treated with palliative systemic chemotherapy and thereafter Palliative radiation. Now she is having stable disease and is on metronomic chemotherapy with regular follow up.

Conclusion: Our case has highlighted that cutaneous metastasis can be seen in the cancers of floor of mouth. Palliative chemotherapy and radiotherapy can significantly increase Progression free survival and hence the Quality of life of patients with such presentation.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Head and Neck cancers account for 7th most common malignancy worldwide. Cutaneous metastasis from head and neck cancers are uncommon.¹ The incidence of cutaneous metastasis from head and neck cancers is less than 1%.² Carcinoma larynx is most often associated out of them.³ It is even rare in cancers of the oral cavity. The common sites of distant metastasis cancers of oral cavity are the lung, liver, and bones.⁴ Here we present a case of carcinoma of floor of mouth with upfront cutaneous metastasis (T4a N2c M1 -Stage 4) treated with Palliative Chemotherapy and Radiotherapy.

2. Case Presentation

A 55-year-old lady presented to our department with complaints of non-healing ulcer oral cavity and multiple swellings over neck from past 6 months. She had a history of local injury after which an ulcer was formed. She had been taking home remedies with no relief. She had no history of any previous radiation treatment, any immunosuppressive agents, drug abuse, tooth extraction or hot beverages.

On examination mouth opening and tongue mobility was restricted with grade 2 trismus. An ulcer proliferative growth approximately of size 3*2 in left GB Sulcus was seen involving the left buccal mucosa. Multiple Bilateral hard fixed lymphadenopathy and multiple hard ulcerated skin nodules on neck and chest wall were noted (Figure 1).

* Corresponding author.

E-mail address: drgrover66@yahoo.in (R. K. Grover).



Figure 1: Cutaneous mets on chest wall



Figure 2: Cutaneous mets on face and neck




Figure 3: Contrast enhanced computed tomography scan of the patient showing cutaneous metastasis.




Figure 4: Fine needle aspiration cytology cutaneous metastasis showing metastatic deposits (Squamous cell carcinoma)

CECT Face and Neck showed heterogeneously enhancing mass 6*5.1cm at the floor of mouth involving left buccal mucosa and erosion of mandible, bilateral lymph node mets 12 to 5 cm and multiple skin nodules over neck and upper chest (Figure 2).

Histopathology of floor of mouth as well as skin nodules were suggestive of moderately differentiated squamous cell carcinoma (Figure 3).

ECOGPS was 1. She was treated with palliative systemic chemotherapy and thereafter Palliative radiation in view of locally advanced disease. Now She is having stable disease on metronomic chemotherapy and is on regular follow up.

3. Discussion

The overall incidence of Cutaneous Metastasis from underlying internal malignancies ranges between 0.7% and 9%, but that from HNSCC is rare (0.7%– 2.4%) with only few cases reported in the literature.⁵ Cutaneous metastasis are defined as isolated or multiple intradermal collections of tumor cells remote from the primary or loco-regional disease.⁶ It is important to differentiate these lesions from extensions from the primary cancer, scar metastasis, metastasis from other primary skin cancers and primary cutaneous disorders.⁷ There are three possible modes of spread namely direct spread, local spread and distant spread. Skin metastasis is thought to involve hematogenous spread where pulmonary circulation and filtration can be theoretically bypassed via the azygous, venous and vertebral venous system via Batson's plexus, allowing skin implantation.⁸

Berger and Fletcher in their study reported that average survival was only around 3 months after skin metastasis becomes clinically evident in HNSCC.⁹ The

treatment of cutaneous metastasis is, in general, palliative. Options available are surgical excision, radiotherapy, and chemotherapy or combination of these. As for now, all patients do poorly and succumb rapidly to the disease.^{10,11} Prakash et al.¹² reported a case of a 78-year-old male with carcinoma of the right buccal mucosa post surgical treatment presented with a near local site recurrence and subsequently developed distant skin metastases to the lower neck and upper trunk during treatment. Verma et al.¹³ also reported a biopsy confirmed cutaneous metastasis from ca tongue post treatment with Surgery and CTRT. Bedi et al.^[14] also reported a case of radically treated case of Ca tongue who presented later on with cutaneous metastasis, treated with palliative chemotherapy and expired 4 months after diagnosis of cutaneous mets. The patient in the present study was treated with palliative chemotherapy and palliative radiation. Now patient is on Metronomic Chemotherapy with stable disease and regular follow up. The patient has been on treatment for more than 14 months. At present, it is difficult to comment on overall survival of the patient.

4. Conclusion

Cutaneous metastasis is a rare presentation in head and neck cancers. Proper investigations and workup should be done for appropriate treatment in such cases. Our case has highlighted that cutaneous metastasis can be seen in the cancers of floor of mouth. Palliative Chemotherapy and Radiotherapy can significantly increase Progression free survival and hence the Quality of life of patients with such presentation.

5. Source of Funding

None.

6. Conflict of Interest


None.

References

1. Krathen RA, Orengo IF, Rosen T. Cutaneous metastasis: a meta-analysis of data. *South Med J*. 2003;96(2):164–7.

2. Pitman KT, Johnson JT. Skin metastases from head and neck squamous cell carcinoma: Incidence and impact. *Head Neck*. 1999;21(6):560–5.
3. Khoury J, Khalifeh I, Kibbi AG, Abbas O. Cutaneous metastasis: Clinicopathological study of 72 patients from a tertiary care center in Lebanon. *Int J Dermatol*. 2014;53(2):147–58.
4. Cooper JS. Carcinoma of the oral cavity and oropharynx. In: Cox J, editor. *Moss' Radiation Oncology: Rationale, Technique, Results*. 7th edn. St. Louis: Mosby; 1994. p. 169–213.
5. Spencer PS, Helm TN. Skin metastases in cancer patients. *Cutis*. 1987;39(2):119–21.
6. Yoskovitch A, Hier MP, Okrainec A, Black MJ, Rochon L. Skin metastases in squamous cell carcinoma of the head and neck. *Otolaryngol Head Neck Surg*. 2001;124(3):248–52.
7. Berger DS, Fletcher GH. Distant metastases following local control of squamous-cell carcinoma of the nasopharynx, tonsillar fossa, and base of the tongue. *Radiology*. 1971;100(1):141–3.
8. Kmucha ST, Troxel JM. Dermal metastases in epidermoid carcinoma of the head and neck. *Arch Otolaryngol Head Neck Surg*. 1993;119(3):326–30.
9. Berger DS, Fletcher GH. Distant metastases following local control of squamous-cell carcinoma of the nasopharynx, tonsillar fossa, and base of the tongue. *Radiology*. 1971;100(1):141–3.
10. Noguti J, Moura CFG, Jesus GPP, Silva VHP, Hossaka TA, Oshima CTF. Metastasis from oral cancer: An overview. *Cancer Genomics Proteomics*. 2012;9(5):329–35.
11. Marioni G, Doro D, Marino F, Verdecchia P, Staffieri C, Staffieri A, et al. Skin and eye: uncommon sites of distant metastasis from tongue base squamous cell carcinoma. *Acta Otolaryngol*. 2003;123(9):1110–4.
12. Prakash A, Upadhyay A. Cutaneous Metastasis of Carcinoma Buccal Mucosa: A Rare Presentation. *Cureus*. 2022;14(6):e25812. doi:10.7759/cureus.25812.
13. Verma K, Gupta M, Gulati A, Sharma RK, Bedi N, Garg P, et al. Multiple cutaneous metastases from squamous cell carcinoma base of tongue: A very rare case report. *J Dermatol Dermatol Surg*. 2016;23(12):858–63.

Author biography

Sheenu Priya, Senior Resident  <https://orcid.org/0009-0000-7272-3548>

Romi Kant Grover, Assistant Professor

Pardeep Garg, Associate Professor

Navik Goyal, Senior Resident

Cite this article: Priya S, Grover RK, Garg P, Goyal N. Cutaneous metastasis in a case of carcinoma floor of mouth- A rare case report. *IP Int J Med Paediatr Oncol* 2024;9(4):147-149.