



Review Article

Orofacial clefts in children and its management

Shaik Ali Hassan^{1,*}, Sumit Bhateja¹, Geetika Arora², Francis Prathyusha³

¹Dept. of Dental, Manav Rachna Dental College, Faridabad, Haryana, India

²Inderprastha Dental College & Hospital, Ghaziabad, Uttar Pradesh, India

³Dr. Francis maxillofacial and dental clinic, India



ARTICLE INFO

Article history:

Received 01-06-2020

Accepted 05-06-2020

Available online 18-07-2020

Keywords:

Orofacial cleft

Zinc

thiopentone

Psychological

Hereditary

ABSTRACT

Orofacial cleft is one of the commonest congenital irregularities which impacts adversely on the life of the individual and to an enormous degree influences the family. Brought about by the collaboration of ecological and hereditary variables, this variation from the norm realizes diminished personal satisfaction. The board of this irregularity involves a group including a split specialist, language teacher, dental specialist, orthodontists, etc. In this review article we will learn about causes, risk factors and various treatment.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Orofacial clefts (OFCs) are common congenital malformations of the lip, sense of taste, or both brought about by complex hereditary and natural components.¹ Orofacial clefts may include the lip, the top of the mouth (hard sense of taste), or the delicate tissue in the rear of the mouth (delicate sense of taste). OFC likewise includes structures around the oral cavity which can broaden onto the facial structures bringing about oral, facial, and craniofacial deformation.² A congenital fissure/sense of taste may affect contrarily on a person's confidence, social abilities, and conduct particularly among young girls.^{3,4} For the most part, young men are influenced more than young ladies with a proportion of around 3:2.⁵ Guys are more probable than females to have a congenital fissure with or without congenital fissure, while females are at a somewhat more serious hazard for congenital fissure alone.^{6,7} Since facial mesenchyme is gotten from neural peak, it is proposed that periconceptual folic corrosive

supplementation may lessen the event of posterity with orofacial clefts.⁸ Zinc additionally is significant in fetal turn of events, and insufficiency of this supplement causes disconnected congenital fissure and other malformations in creatures; different supplements, for example, riboflavin and nutrient and are additionally fundamental.⁹ Preventive endeavors may involve control of maternal way of life, improved eating routine and utilization of multivitamin and mineral enhancements, evasion of certain medications and drugs, and general awareness of social, word related, and private hazard factors.

1.1. Genetic basis of clefts

Hereditary legacy implies that a kid's highlights are "acquired" or went from parent to kid.¹⁰ There are two kinds of legacy: the single quality legacy where an element shows up because of a single quality conveyed by one parent and the multifactorial legacy where an element shows up because of a number of hereditary and nongenetic factors, for example, liquor, drugs, also, natural variables.¹⁰ Orofacial advancement is a mind-boggling process that

* Corresponding author.

E-mail address: alishaikhassan@gmail.com (S. A. Hassan).

includes numerous qualities and flagging pathways.¹¹

1.2. Risk of occurrence

The danger of repeat of a separated condition is controlled by various components that are regularly exceptional in a specific family, and these incorporate the quantity of relatives with clefts, how firmly related these individuals are, the race and sex of the influenced people, and the sort of parted every individual has.¹² When parents have a youngster with a parted, the hazard that the following youngster (each succeeding kid) will be influenced is 2–5% (2 to 5 possibilities in 100). In the event that there is more than one individual in the close family with a separated, the hazard ascends to 10–12% (about 1 possibility in 10). A person who is the just one in their family with a parted has 2–5% chances that their kid will have a parted (2 to 5 possibilities in 100). On the off chance that the person with the separated likewise has a nearby family member with a parted, the hazard increments to 10–12% (around 1 out of 10) that a kid will have a separated.¹³ At last, the unaffected kind of a person with a parted have a generally 1% (1 of every 100) chance of having a child with a separated. This may ascend to 5–6% (5 to 6 risks in 100) if more than one close relative has a split.

1.3. Causes for cleft

The etiology of OFCs is unpredictable, including various hereditary and natural elements.¹⁴ Oral clefts much of the time happen in mix with a wide scope of chromosomal variations from the norm and conditions (trisomy 13, amniotic band anomalad, Fryns disorder, Meckel condition, Fanatic condition, Treacher Collins disorder, van der Woude condition, Velocardiofacial disorder, and so on.)¹⁵ and natural factors, for example, drug during pregnancy, maternal liquor utilization and smoking, dietary and nutrient lacks, diabetes, ecological poisons, elevation, birth request, financial status, and parental age.^{16–20} Other hereditary variables that may influence the nearness of OFCs incorporate maternal capacity to keep up red platelet zinc focuses and myoinositol fixations (a hexahydroxycyclohexane sugar liquor).²¹ Maternal capacity to keep up sufficient degrees of nutrients B6 and B12 and fetal capacity to use these supplements are additionally observed as a factor in the advancement of oral clefts when these supplements are not used appropriately since mistakes in DNA blend and interpretation may happen.²²

Medications assume a constrained job in the etiology of congenital fissure or then again sense of taste (CL/P); amoxicillin, phenytoin, oxprenolol, and thiethylperazine may have some relationship with (CL/P), while carbamazepine and oxytetracycline may have a few relationship with back congenital fissure (PCP) during the beginning periods of pregnancy (23). What's more, moms with the

MTHFR 677TT OR MTHFR 1298CC genotype and low periconceptional folate admission were found to have an expanded hazard for congenital fissure with or without sense of taste among their posterity.^{23,24} Medications which meddle with folate digestion, for example, phenytoin, are known to have teratogenic impacts which incorporate oral split, development hindrance, appendage abandons, and other craniofacial distortions. Maternal admission of vasoactive medications which incorporate pseudoephedrine, headache medicine, ibuprofen, and amphetamine just as cigarette smoking has been related with higher danger of oral clefts.^{25–27}

1.4. Diagnosis

In some cases OFCs are analyzed by pre-birth ultrasound, yet there is no precise screening for orofacial clefts.²⁸ Regularly, orofacial clefts are analyzed after the child is conceived. Nonetheless, here and there minor clefts (e.g., submucous congenital fissure and bifid uvula) probably won't be analyzed until further down the road. Congenital fissure can be handily analyzed by performing ultrasonography in the second trimester of pregnancy when the situation of the fetal face is found effectively.²⁹

1.5. Latest diagnosis ways

New methods for pre-birth finding have been accounted for by certain creators. Campbell et al. announced a novel strategy including a converse face see utilizing 3D sonography to analyze congenital fissure and sense of taste in the antenatal period.^{30,31} While pre-birth finding of parted is promptly feasible utilizing regular 2D sonography, congenital fissure is increasingly hard to distinguish particularly on the off chance that it is an related inconsistency.³² Platt et al. accepted that the exact analysis of craniofacial malformations can be upgraded with 3D sonography.³³ Spoiled and Levaillant (2004) have likewise announced that incorporation of 3D and 4D ultrasound imaging permits simpler and increasingly exact assessment of the distinctive separated constituents.³⁴

1.6. Problem with treatment

Treatment of congenital fissure and sense of taste irregularities requires long periods of particular care and is expensive. The normal lifetime clinical cost for treatment of one individual influenced with a congenital fissure, what's more, sense of taste is \$100,000.³⁵ Albeit fruitful treatment of the restorative and utilitarian parts of orofacial separated oddities is currently conceivable, it is as yet testing, extensive, exorbitant, and reliant on the aptitudes and experience of a clinical group. This particularly applies to careful, dental, furthermore, language courses.³⁶ The mean and middle expenses for youngsters ≤10 years old with an orofacial separated were eight times higher than

those for offspring of a similar age without an orofacial separated. Mean expenses for newborn children with a split and another major irrelevant imperfection were multiple times higher than those for a baby without a split and multiple times higher than those for babies with a secluded split with patients consistently joined up with a charge for-administration.³⁷ New methods for pre-birth conclusion have been accounted for by certain creators. Campbell et al. detailed a novel method including a converse face see utilizing 3D sonography to analyze congenital fissure and sense of taste in the antenatal period.^{38,39} While pre-birth finding of parted is promptly achievable utilizing regular 2D sonography, congenital fissure is progressively hard to distinguish particularly in the event that it is an related inconsistency.⁴⁰ Platt et al. accepted that the precise finding of craniofacial malformations can be upgraded with 3D sonography.⁴¹ Spoiled and Levailant (2004) have additionally revealed that consideration of 3D and 4D ultrasound imaging permits simpler and progressively exact assessment of the diverse parted constituents.⁴²

2. Management

2.1. Sugical management

Orofacial clefts by and large require careful fix. Regularly different medical procedures are expected to remake the lip and sense of taste.⁴³ A palatoplasty is the method used to close the sense of taste, reestablish the velopharyngeal sphincter, and assist discourse with working and different procedures.⁴⁴ The ideal way to deal with the treatment of youngsters brought into the world with parted deformities is a multidisciplinary approach which includes joined endeavors of a pediatrician, orthodontist, master nurture, separated specialist, language teacher, and ear, nose, what's more, throat pro to give the best consolidated ability to guarantee that the right intercessions are done at the suitable time and to guarantee the best useful and stylish outcome.⁴⁴ Numerous youngsters will require extra medical procedures as they get more seasoned. Careful fix can improve the look and presence of a youngster's face, it likewise may improve breathing, and shearing, discourse and language.⁴⁵

2.2. Medical management

The supplementation of folic corrosive at present prescribed to secure against neural cylinder deserts is 0.4 mg every day, double the present normal day by day consumption for ladies of 0.2 mg.⁴⁶ It has been proposed that maternal folic corrosive supplementation assumes a job in the counteraction of non-syndromic orofacial clefts, that is, congenital fissure with or without congenital fissure (CL ± P) (48). A few investigations have revealed diminished paces of congenital fissure and sense of taste with folic corrosive use.^{47–50} Some vagueness of the investigations might be clarified by an ongoing report that found that

oral split hazard can be decreased just by high portions of folic corrosive devoured at the hour of lip furthermore, sense of taste development.⁵¹ Maternal multivitamin use has additionally been found to bring about a noteworthy decrease in split sense of taste chance and a nonsignificant decrease in congenital fissure hazard.⁵²

2.3. Psychological management

The mental care of the understanding with a parted starts at the hour of determination, regardless of whether this is before birth. An exact analysis is basic to the procedure of guiding families. It is the duty of the referral focus to characterize the idea of the auxiliary deformity with as much exactness as could be expected under the circumstances. This causes the family to envision the kid and to talk about taking care of, particularly breastfeeding. It additionally helps when illuminating about planning and sort of medical procedure. To get ready for the future, parents need to talk about the executives and likely the treatment pathway at their own pace and at their own time, with the goal that they are ready to retain the data.⁵³ Postponed fix of split can prompt debilitated family and cultural associations with potential long haul mental consequences for the kid. As the kid develops and faces the errand of individuation from the family, there might be a requirement for mental work, and since adulthood gives its own arrangement of difficulties to the person, there is potential for further mental intercessions all through this time of life.

2.4. Other treatment options

For different medicines such as hearing appraisal, discourse and language treatment, and dentofacial advancement and treatment, therapist or other mental authorities are required to guarantee powerful working of the body organs and frameworks.⁵⁴ The job of craniofacial group in the administration of congenital fissure ± sense of taste can't be downplayed. A craniofacial group is a multidisciplinary group which gives multidisciplinary interviews, finding, treatment arranging, and methods for a range of craniofacial abnormalities and conditions.⁵⁵ Cooperation is strongly suggested in the administration of people with OFC. This group is highly devoted to guaranteeing that people with the condition are offered the vital assistance, care, and backing to assist them with having a superior life.

3. Conclusion

Instruction and awareness on orofacial clefts all in all ought to be advanced so preventive measures can be set up what's more, people experiencing the condition can be very much joined in to and provided food for. Orofacial clefts sway on an individual's personal satisfaction subsequently the requirement for better administration of this variation from the norm. There is a requirement for additional

examinations to be conveyed out on split hereditary qualities.

4. Source of Funding

None.

5. Conflict of Interest

None.

References

- Wehby GL, Murray JC. Folic acid and orofacial clefts: a review of the evidence. *Oral Dis.* 2010;16(1):11–9.
- Mossey P, Little J. Addressing the challenges of cleft lip and palate research in India. *Indian J Plast Surg.* 2009;42(1):S9–18.
- Cleft Lip and Palate. Available from: <http://en.wikipedia.org/wiki/Cleftlipandpalate>.
- Leonard BJ, Brust JD. Self-concept of children and adolescents with cleft lip and/or palate. *Cleft Palate-Craniofacial J.* 1991;28(4):347–53.
- Ellis E. Management of patients with orofacial clefts,” in Contemporary Oral and Maxillofacial Surgery. Mosby, St. Louis, Mo, USA; 2003. p. 623–45.
- Blanco-Davila F. Incidence of Cleft Lip and Palate in the Northeast of Mexico: A 10-Year Study. *J Craniofac Surg.* 2003;14(4):533–7.
- Gregg T, Boyd D, Richardson A. The Incidence of Cleft Lip and Palate in Northern Ireland from 1980–1990. *Br J Orthod.* 1994;21(4):387–92.
- Hartridge T, Illing HM, Sandy JR. The Role of Folic Acid in Oral Clefting. *Br J Orthod.* 1999;26(2):115–20.
- Rothman KJ, Moore LL, Singer MR, Nguyen USDT, Mannino S, Milunsky A. Teratogenicity of High Vitamin A Intake. *N Engl J Med.* 1995;333(21):1369–73.
- White Memorial Medical Center (Adventist Health) Cleft Palate Program; Available from: <http://www.whitememorial.com/medical-services/cleft-palate-faq>.
- Murray JC, Schutte BC. Cleft palate: players, pathways, and pursuits. *J Clin Invest.* 2004;113(12):1676–8.
- Cleft palate Foundation; Available from: <http://www.cleftline.org/docs/Booklets/GEN-01.pdf>.
- Agbenorku P, Agbenorku M, Iddi A, Abude F, Sefenu R, Matondo P, et al. A study of cleft lip/palate in a community in the South East of Ghana. *Eur J Plast Surg.* 2011;34(4):267–72.
- Schutte BC. The many faces and factors of orofacial clefts. *Human Mol Genet.* 1999;8(10):1853–9.
- Birth Defect Risk Factor Series: Oral Clefts; 2012.
- Little J, Cardy A, Munger RG. Tobacco smoking and oral clefts: a meta-analysis. *Bull World Health Organ.* 2004;82(3):213–8.
- Spilson SV, Kim HJE, Chung KC. Association Between Maternal Diabetes Mellitus and Newborn Oral Cleft. *Ann Plast Surg.* 2001;47(5):477–81.
- Castilla EE, Lopez-Camelo JS, Campaa H. Altitude as a risk factor for congenital anomalies. *Am J Med Genet.* 1999;86(1):9–14.
- Vieira AR, Orioli IM, Murray JC. Maternal age and oral clefts: a reappraisal,” Oral Surgery, Oral Medicine, Oral Pathology. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol, Endod.* 2002;94(5):530–5.
- Yang J, Carmichael SL, Canfield M, Song J, and GMS. Socioeconomic Status in Relation to Selected Birth Defects in a Large Multicentered US Case-Control Study. *Am J Epidemiol.* 2007;167(2):145–54.
- Krapels IPC, Rooij IALM, Wevers RA, Zielhuis GA, Spauwen PHM, Brussel W, et al. Myo-inositol, glucose and zinc status as risk factors for non-syndromic cleft lip with or without cleft palate in offspring: a case-control study. *BJOG: An Int J Obstet Gynaecol.* 2004;111(7):661–8.
- van Rooij IALM. Does the Interaction between Maternal Folate Intake and the Methylenetetrahydrofolate Reductase Polymorphisms Affect the Risk of Cleft Lip with or without Cleft Palate? *Am J Epidemiol.* 2003;157(7):583–91.
- Msamati BC, Igbigbi PS, Chisi JE. The incidence of cleft lip, cleft palate, hydrocephalus and <i>spina bifida</i> at Queen Elizabeth Central Hospital, Blantyre, Malawi. *Cent Afr J Med.* 2000;46(11):292–6.
- Beaty TH, Maestri NE, Hetmanski JB. Testing for interaction between maternal smoking and TGFA genotype among oral cleft cases born in Maryland 1992–1996. *Cleft Palate-Craniofacial J.* 1997;34:447–54.
- Lammer EJ, Shaw GM, Iovannisci DM, Finnell RH. Periconceptional multivitamin intake during early pregnancy, genetic variation of acetyl-N-transferase 1 (NAT1), and risk for orofacial clefts. *Birth Defects Res Clin Mol Teratology.* 2004;70(11):846–52.
- Munger RG, Romitti PA, Daack-Hirsch S, Burns TL, Murray JC, Hanson J, et al. Maternal alcohol use and risk of orofacial cleft birth defects. *Teratology.* 1996;54(1):27–33.
- Utah Department of Health: Striving to Prevent Birth Defects. *Orofacial Clefts at a Glance*.
- Centers for Disease Control and Prevention, Birth Defects: Facts about Cleft Lip and Cleft Palate; Available from: <http://www.cdc.gov/ncbddd/birthdefects/cleftlip.html>.
- Kall B. Maternal drug use and infant cleft lip/palate with special reference to corticoids. *Cleft Palate-Craniofac J.* 2003;40(6):624–8.
- Campbell S, Lees CC. The three-dimensional reverse face (3D RF) view for the diagnosis of cleft palate. *Ultrasound Obstet Gynecol.* 2003;22(5):552–4.
- Campbell S, Lees C, Moscoso G, Hall P. Ultrasound antenatal diagnosis of cleft palate by a new technique: the 3D ‘reverse face’ view. *Ultrasound Obstet Gynecol.* 2005;25(1):12–8.
- Walker SJ, Ball RH, Babcock CJ, Feldkamp MM. Prevalence of aneuploidy and additional anatomic abnormalities in fetuses and neonates with cleft lip with or without cleft palate: a population-based study in Utah. *J Ultrasound Med.* 2001;20(11):1175–80.
- Platt LD, DeVore GR, Pretorius DH. Improving Cleft Palate/Cleft Lip Antenatal Diagnosis by 3-Dimensional Sonography. *J Ultrasound Med.* 2006;25(11):1423–30.
- Rotten D, Levaillant JM. Two- and three-dimensional sonographic assessment of the fetal face. 2. Analysis of cleft lip, alveolus and palate. *Ultrasound Obstet Gynecol.* 2004;24(4):402–11.
- Waitzman NJ, Romano PS, Scheffler RM. Estimates of the economic costs of birth defects. *Inquiry.* 1994;31(2):188–205.
- Centers for Disease Control and Prevention, Birth Defects: Facts about Cleft Lip and Cleft Palate; Available from: <http://www.cdc.gov/ncbddd/birthdefects/cleftlip.html>.
- Boulet SL, Grosse SD, Honein MA, Correa-Villaseñor A. Children with Orofacial Clefts: Health-Care Use and Costs among a Privately Insured Population. *Public Health Rep.* 2009;124(3):447–53.
- Quaderi NA, Schweiger S, Gaudenz K, Franco B, Rugarli EI, Berger W, et al. Opitz G/BBB syndrome, a defect of midline development, is due to mutations in a new RING finger gene on Xp22. *Nat Genet.* 1997;17(3):285–91.
- Rijli FM, Mark M, Lakkaraju S, Dierich A, Dollé P, Chambon P, et al. A homeotic transformation is generated in the rostral branchial region of the head by disruption of Hoxa-2, which acts as a selector gene. *Cell.* 1993;75(7):1333–49.
- Satokata I, Maas R. Msx1 deficient mice exhibit cleft palate and abnormalities of craniofacial and tooth development. *Nat Genet.* 1994;6(4):348–56.
- Zhao Y, Guo YJ, Tomac AC, Taylor NR, Grinberg A, Lee EJ, et al. Isolated cleft palate in mice with a targeted mutation of the LIM homeobox gene Lhx8. *Proc National Acad Sci.* 1999;96(26):15002–6.
- Waitzman NJ, Romano PS, Scheffler RM. Estimates of the economic costs of birth defects. *Inquiry.* 1994;31(2):188–205.
- Werler MM, Hayes C, Louik C, Shapiro S, Mitchell AA. Multivitamin Supplementation and Risk of Birth Defects. *Am J Epidemiol.* 1999;150(7):675–82.
- Brailsford J, Smith DD, Lizarraga AK, Bermudez LE. Surgical management of patients with cleft palate. *OR Nurse.* 2010;4(3):16–

- 25.
45. Mossey PA, Munger R, Murray JC, Shaw WC. WHO Reports, Human Genetics Programme, "Management of non-communicable diseases. International collaborative research on craniofacial anomalies," in Global Strategies Towards Reducing the Health Care Burden of Craniofacial Anomalies. 2002;.
 46. Menulty H. Folate requirements for health in different population groups. *Br J Biomed Sci.* 1995;52(2):110–9.
 47. Malek FA, Möritz KU, Fanghänel J, Bienengraber V. Sex-related differences in procarbazine-induced cleft palate and microgenia and the anti-teratogenic effect of prenatal folic acid supplementation in rats. *Ann Anat.* 2003;185(5):465–70.
 48. Mulinare J, Erickson JD, James LM. Does periconceptional use of multivitamins reduce the occurrence of birth defects? *Am J Epidemiol.* 1995;141(3).
 49. Munger R, Romitti P, West N. Maternal intake of folate, vitamin B-12, and zinc and risk of orofacial cleft birth defects. *Am J Epidemiol.* 1997;145(30).
 50. Shaw GM, Nelson V, Carmichael SL, Lammer EJ, Finnell RH, Rosenquist TH, et al. Maternal Periconceptional Vitamins: Interactions with Selected Factors and Congenital Anomalies? *Epidemiol.* 2002;13(6):625–30.
 51. Czeizel AE, Timar L, Sarkozi A. Dose-dependent Effect of Folic Acid on the Prevention of Orofacial Clefts. *Pediatr.* 1999;104(6):e66.
 52. Werler MM, Hayes C, Louik C, Shapiro S, Mitchell AA. Multivitamin Supplementation and Risk of Brith Defects. *Am J Epidemiol.* 1999;150(7):675–82.
 53. Hodgkinson PD, Brown S, Duncan D. Management of children with cleft lip and palate: a review describing the application of multidisciplinary team working in this condition based upon the experiences of a regional cleft lip and palate centre in the United Kingdom. *Fetal Matern Med Rev.* 2005;16(1):1–27.
 54. Bristow L, Bristow S. Making Faces: Logan's Cleft Lip and Palate Story, Pulsus Group, Ontario, Canada. 2007;.
 55. Tolarova MM, Poulton D, Aubert MM. Pacific ´craniofacial team and cleft prevention program. *Journal of the California Dental Association.* 2006;34(10):823–830.

Author biography

Shaik Ali Hassan Dental Surgeon

Sumit Bhateja HOD

Geetika Arora Reader

Francis Prathyusha BDS, MDS

Cite this article: Hassan SA, Bhateja S, Arora G, Prathyusha F. **Orofacial clefts in children and its management.** *IP Int J Med Paediatr Oncol* 2020;6(2):38-42.